Index of Authors

ADAMS, P. A. The kinetics of the recombination reaction between apomyoglobin and alkaline haematin 153–158

ÅGREN, A. *see* BROLIN, S. E. 159–162

ALDRIDGE, W. N. *see* ELLIOTT, B. M. 583–589

ALLEN, A. *see* SCAWEN, M. 363–368

ALLEN, G. *see* BAINES, B. S. 189–192

ANDERSSON, S. M. *see* OHISALO, J. J. 411–417

AUSTEN, B. M. & MARSHALL, R. D. Acid dissociation constants of glycopeptides 31–38

AXEL'ROD, V. D. *see* GORBULEV, V. G. 409–410

BAILEY, A. J. *see* ROBINS, S. P. 339–346

BAINES, B. S., ALLEN, G. & BROCKLEHURST, K. The highly electrophilic character of 4-chloro-7-nitrobenzofurazan and possible consequences for its applications as a protein-labelling reagent 189–192

BAIRD, M. B., MASSIE, H. R. & BIRNBAUM, L. S. Presence of a high-molecular-weight form of catalase in enzyme purified from mouse liver 449–453

BANDURSKI, R. S. *see* DILWORTH, G. L. 521–529

BARBER, D., PARR, S. R. & GREENWOOD, C. The reduction of *Pseudomonas* cytochrome c_{551} oxidase by chromous ions 629–632

BARBER, M. J., BRAY, R. C., CAMMACK, R. & COUGHLAN, M. P. Oxidation–reduction potentials of turkey liver xanthine dehydrogenase and the origins of oxidase and dehydrogenase behaviour in molybdenum-containing hydroxylases 279–289

BARCISZEWSKI, J. *see* GORBULEV, V. G. 409–410

BAYLEY, P. M. *see* CHEN, S.-S. 297–302

BEARD, J. *see* SIMPSON, I. A. 309–316

BEAVEN, G. H. *see* GEISOW, M. J. 477–484

BELL, M. V., TONDEUR, F. & SARGENT, J. R. The activation of sodium-plus-potassium ion-dependent adenosine triphosphatase from marine teleost gills by univalent cations 185–187

BELL-BOOTH, P. G. *see* GILLARD, G. C. 145–151

BENNICK, A. Chemical and physical characterization of a phosphoprotein, protein C, from human saliva and comparison with a related protein A 229–239

BENNICK, A. The binding of calcium to salivary phosphoprotein, protein C, and comparison with calcium binding to protein A, a related salivary phosphoprotein 241–245

BENSADOUN, A. *see* EHNHOLM, C. 347–355

BERTI, J. D. *see* CICILINI, M. A. 433–439

BEVAN, M. J. *see* TROWBRIDGE, I. S. 211–217

BIRNBAUM, L. S. *see* BAIRD, M. B. 449–453

BISHOP, S. H. *see* GIBBS, K. L. 511–516

BLOXHAM, D. P. *see* RAGAN, C. I. 605–615

BOHM, L. D. *see* HOWLAND, R. D. 125–131

BOOTH, A. G. A novel system for the two-dimensional electrophoresis of membrane proteins 165–168

BOULTER, D. *see* LYDDIATT, A. 333–338

BRATTAIN, M. G., KIMBALL, P. M., PRETLOW, T. G., II & MARKS, M. E. The interaction of *N*-acetylhexosaminidase with insolubilized concanavalin A 247–251

BRAY, R. C. *see* BARBER, M. J. 279–289

BRESCIANI, D. Different reactivities of free and bound amino groups in deoxy- and liganded haemoglobin 393–395

BRILEY, P. A., EISENTHAL, R., HARRISON, R. & SMITH, G. D. [^{19}F]Fluorine nuclear-magnetic-resonance study of the interaction of difluoro-oxaloacetate with aspartate transaminase 325–331

BROCKLEHURST, K. Evolution of enzyme catalytic power. Characteristics of optimal catalysis evaluated for the simplest plausible kinetic model 111–116

BROCKLEHURST, K. *see also* BAINES, B. S. 189–192

BROLIN, S. E. & ÅGREN, A. Assay of flavin nucleotides in pancreatic islets by a differential fluorimetric technique 159–162

BROWN, A. S. & TROXLER, R. F. Properties and *N*-terminal sequence of allophycocyanin from the unicellular rhodophyte *Cyanidium caldarium* 571–581

BROWN, F., NEAL, D. J. & WILKINSON, S. G. Presence of 3-*O*-methyl-L-xylose in the lipopolysaccharide of *Pseudomonas maltophilia* N.C.T.C. 10257 173–175

BROWN, W. V. *see* EHNHOLM, C. 347–355

BURCHELL, B. *see* KNOWLES, R. G. 381–383

BURY, A. F., COOLBEAR, T. & SAVERY, C. R. Separation and properties of two arylamidases from rat cardiac-muscle extracts 565–570

CALDO, H. *see* CICILINI, M. A. 433–439

CAMARGO, A. C. M. *see* CICILINI, M. A. 433–439

CAMMACK, R. *see* BARBER, M. J. 279–289

CAMPBELL, A. M. & LOWE, P. A. The dissociation of σ-factor from ribonucleic acid polymerase 177–179

CARTWRIGHT, E. *see* SELLERS, A. 303–307

CHANG, J. Y. High-sensitivity sequence analysis of peptides and proteins by 4-*NN*-dimethylaminoazobenzene 4′-isothiocyanate 517–520

CHAPMAN, M. J. *see* MILLS, G. L. 455–465

CHEAH, K. S. E. & OSBORNE, D. J. Analysis of nucleosomal deoxyribonucleic acid in a higher plant 141–144

CHEN, S.-S., ENGEL, P. C. & BAYLEY, P. M. The allosteric mechanism of bovine liver glutamate dehydrogenase. Evidence from circular-dichroism studies for a conformational change in the ternary complex enzyme–(oxidized nicotanimide–adenine dinucleotide)–glutarate 297–302

CHRISTENSEN, U. & CLEMMENSEN, I. Kinetic properties of the primary inhibitor of plasmin from human plasma 389–391

CICILINI, M. A., CALDO, H., BERTI, J. D. & CAMARGO, A. C. M. Rabbit tissue peptidases that hydrolyse the peptide hormone bradykinin. Differences in enzymic properties and concentration in rabbit tissues 433–439

CLEMMENSEN, I. *see* CHRISTENSEN, U. 389–391

COOLBEAR, T. *see* BURY, A. F. 565–570

COUGHLAN, M. P. *see* BARBER, M. J. 279–289

CRISSEY, M. H. *see* VESSEY, D. A. 181–183

CROISSANT, R. D. *see* GUENTHER, H. L. 591–603

DICKINSON, F. M. *see* HART, G. J. 261–267

DILWORTH, G. L. & BANDURSKI, R. S. Activation of selenate by adenosine 5′-triphosphate sulphurylase from *Saccharomyces cerevisiae* 521–529

DUGAL, B. Measurement of 1-aspartamido-β-*N*-acetylglucosamine amidohydrolase activity in human tissues 9–14

DUNKER, A. K. & ZALESKE, D. J. Stereochemical considerations for constructing α-helical protein bundles with particular application to membrane proteins 45–57
DUNKLEY, P. R., HOLMES, H. & RODNIGHT, R. Phosphorylation of synaptic-membrane proteins from ox cerebral cortex *in vitro*. Preparation of fractions enriched in phosphorylated proteins by using extraction with detergents and urea, and gel filtration 369–378

ECCLESTON, J. F. & TRENTHAM, D. R. The interaction of chromophoric nucleotides with subfragment 1 of myosin 15–29
EHNHOLM, C., BENSADOUN, A. & BROWN, W. V. Characterization of two triacylglycerol lipase activities in pig post-heparin plasma 347–355
EISENTHAL, R. *see* BRILEY, P. A. 325–331
ELLIOTT, B. M. & ALDRIDGE, W. N. Binding of triethyltin to cat haemoglobin and modification of the binding sites by diethyl pyrocarbonate 583–589
ENGEL, P. C. *see* CHEN, S.-S. 297–302
ERDÖS, E. G. *see* MARINKOVIC, D. V. 253–260
ERFLE, J. D. *see* MAHADEVAN, S. 495–501

FLINT, M. H. *see* GILLARD, G. C. 145–151
FORSTER, G. R. *see* MILLS, G. L. 455–465
FRIEDMAN, P. A. & SHIA, M. A. The apparent absence of involvement of biotin in the vitamin K-dependent carboxylation of glutamic acid residues of proteins 39–43

GEDDES, R., HARVEY, J. D. & WILLS, P. R. The molecular size and shape of liver glycogen 201–209
GEDDES, R. & STRATTON, G. C. The influence of lysosomes on glycogen metabolism 193–200
GEISOW, M. J. & BEAVEN, G. H. Physical and binding properties of large fragments of human serum albumin 477–484
GIBBS, K. L. & BISHOP, S. H. Adenosine triphosphate-activated adenylate deaminase from marine invertebrate animals. Properties of the enzyme from lugworm (*Arenicola cristata*) body-wall muscle 511–516
GIGLI, I. *see* SIM, R. B. 219–227
GILLARD, G. C., MERRILEES, M. J., BELL-BOOTH, P. G., REILLY, H. C. & FLINT, M. H. The proteoglycan content and the axial periodicity of collagen in tendon 145–151
GORBULEV, V. G., KUTATELADZE, T. V., BARCISZEWSKI, J. & AXEL'ROD, V. D. The separation of oligonucleotides of baker's-yeast valine transfer ribonucleic acid 2b by high-voltage electrophoresis on DEAE-paper and by thin-layer chromatography 409–410
GOSO, K. & HOTTA, K. A glycopeptide isolated from human gastric juice 169–172
GRAHAM, A. B., PECHEY, D. T., TOOGOOD, K. C., THOMAS, S. B. & WOOD, G. C. The phospholipid-dependence of uridine diphosphate glucuronyltransferase. Phospholipid depletion and re-activation of guinea-pig liver microsomal enzyme 117–124
GRAY, C. H. *see* STOLL, M. S. 59–101
GREENWOOD, C. *see* BARBER, D. 629–632
GUENTHER, H. L., CROISSANT, R. D., SCHONFELD, S. E. & SLAVKIN, H. C. Identification of four extracellular-matrix enamel proteins during embryonic-rabbit tooth-organ development 591–603

HALLIWELL, B. Generation of hydrogen peroxide, superoxide and hydroxyl radicals during the oxidation of dihydroxyfumaric acid by peroxidase 441–448
HAMILTON, R. T. *see* TROWBRIDGE, I. S. 211–217
HARRISON, R. *see* BRILEY, P. A. 325–331
HART, G. J. & DICKINSON, F. M. Some properties of aldehyde dehydrogenase from sheep liver mitochondria 261–267
HARVEY, J. D. *see* GEDDES, R. 201–209
HAVINGA, E. *see* VAN BATENBURG, O. D. 385–387
HOES, K. *see* VAN BATENBURG, O. D. 385–387
HOLLAWAY, M. R. *see* SIMPSON, I. A. 309–316
HOLMES, H. *see* DUNKLEY, P. R. 369–378
HOTTA, K. *see* GOSO, K. 169–172
HOWLAND, R. D. & BOHM, L. D. Possible multiple binding sites for *o*-aminophenol on uridine diphosphate glucuronyltransferase 125–131
HUGHES, S. G. A map of the cleavage sites for endonuclease *Ava*I in the chromosome of bacteriophage lambda 503–509

INDGE, K. J. Application of topological methods in enzyme kinetics 633–634
INTERUNION COMMISSION ON BIOTHERMODYNAMICS. Recommendations for measurement and presentation of biochemical equilibrium data 1–7

JEWETT, P. B. *see* NEUWELT, E. A. 419–426
JOHNSON, D. & TRAVIS, J. Inactivation of human α_1 proteinase inhibitor by thiol proteinases 639–641
JOLLY, R. D. *see* PHILLIPS, N. C. 269–277

KAGEURA, E. & TOKI, S. Purification and properties of a new testosterone 17β-dehydrogenase ($NADP^+$) from guinea-pig liver 401–407
KERLING, K. E. T. *see* VAN BATENBURG, O. D. 385–387
KIMBALL, P. M. *see* BRATTAIN, M. G. 247–251
KNOWLES, R. G. & BURCHELL, B. A simple method for purification of epoxide hydratase from rat liver 381–383
KUTATELADZE, T. V. *see* GORBULEV, V. G. 409–410

LANDT, M. L. *see* SWISHER, R. H. 427–432
LAURENT, T. C. *see* PEARCE, R. H. 617–625
LEHMAN, W. Calcium ion-dependent myosin from decapod-crustacean muscles 291–296
LEVY, C. C. *see* NEUWELT, E. A. 419–426
LONDESBOROUGH, J. Characterization of an adenosine 3′:5′-cyclic monophosphate phosphodiesterase from baker's yeast. Its binding to subcellular particles, catalytic properties and gel-filtration behaviour 467–476
LOWE, P. A. *see* CAMPBELL, A. M. 177–179
LUNT, M. R. *see* ROSAMOND, J. D. C. 485–494
LYDDIATT, A. & BOULTER, D. The amino acid sequence of cytochrome *c* from the locust, *Schistocerca gregaria* Forskal 333–338

MAHADEVAN, S., SAUER, F. D. & ERFLE, J. D. Purification and properties of urease from bovine rumen 495–501
MARINKOVIC, D. V. & MARINKOVIC, J. N. Purification of two hexosaminidases from human kidney 133–140

MARINKOVIC, D. V., MARINKOVIC, J. N., ERDŐS, E. G. & ROBINSON, C. J. G. Purification of carboxypeptidase B from human pancreas 253–260
MARINKOVIC, J. N. *see* MARINKOVIC, D. V. 133–140, 253–260
MARKS, M. E. *see* BRATTAIN, M. G. 247–251
MAROTEAUX, P. *see* STANESCU, V. 103–109
MARSHALL, R. D. *see* AUSTEN, B. M. 31–38
MASSIE, H. R. *see* BAIRD, M. B. 449–453
MERRILEES, M. J. *see* GILLARD, G. C. 145–151
MILLS, G. L., TAYLAUR, C. E., CHAPMAN, M. J. & FORSTER, G. R. Characterization of serum lipoproteins of the shark *Centrophorus squamosus* 455–465
MORIHARA, K. & OKA, T. α-Chymotrypsin as the catalyst for peptide synthesis 531–542
MURPHY, G. *see* SELLERS, A. 303–307

NEAL, D. J. *see* BROWN, F. 173–175
NEUWELT, E. A., SCHMUKLER, M., NIZIAK, M. S., JEWETT, P. B. & LEVY, C. C. The immunological characterization of several human ribonucleases by using primary binding tests 419–426
NILSEN-HAMILTON, M. *see* TROWBRIDGE, I. S. 211–217
NIZIAK, M. S. *see* NEUWELT, E. A. 419–426

OHISALO, J. J., ANDERSSON, S. M. & PISPA, J. P. Partial purification and properties of frog liver tyrosine aminotransferase 411–417
OKA, T. *see* MORIHARA, K. 531–542
OSBORNE, D. J. *see* CHEAH, K. S. E. 141–144

PARR, S. R. *see* BARBER, D. 629–632
PASNAK, C. F. *see* QUARLES, R. H. 635–637
PEARCE, R. H. & LAURENT, T. C. Exclusion of dextrans by meshworks of collagenous fibres 617–625
PECHEY, D. T. *see* GRAHAM, A. B. 117–124
PHILLIPS, N. C., WINCHESTER, B. G. & JOLLY, R. D. A serological investigation into the acidic α-D-mannosidase in normal Angus cattle and in a calf with mannosidosis 269–277
PISPA, J. P. *see* OHISALO, J. J. 411–417
PORTER, R. R. *see* SIM, R. B. 219–227
PRETLOW, T. G., II. *see* BRATTAIN, M. G. 247–251

QUARLES, R. H. & PASNAK, C. F. A rapid procedure for selectively isolating the major glycoprotein from purified rat brain myelin 635–637

RAGAN, C. I. & BLOXHAM, D. P. Specific labelling of a constituent polypeptide of bovine heart mitochondrial reduced nicotinamide–adenine dinucleotide–ubiquinone reductase by the inhibitor diphenyleneiodonium 605–615
REID, K. B. M. *see* SIM, R. B. 219–227
REILLY, H. C. *see* GILLARD, G. C. 145–151
REITHEL, F. J. *see* SWISHER, R. H. 427–432
REYNOLDS, J. J. *see* SELLERS, A. 303–307
ROBINS, S. P. & BAILEY, A. J. The chemistry of the collagen cross-links. Characterization of the products of reduction of skin, tendon and bone with sodium cyanoborohydride 339–346
ROBINSON, C. J. G. *see* MARINKOVIC, D. V. 253–260
ROCHE, J. J. *see* SPRAGG, S. P. 543–555
RODNIGHT, R. *see* DUNKLEY, P. R. 369–378
ROSAMOND, J. D. C. & LUNT, M. R. An adenosine triphosphate-dependent deoxyribonuclease from *Alcaligenes faecalis* 485–494

SARGENT, J. R. *see* BELL, M. V. 185–187
SAUER, F. D. *see* MAHADEVAN, S. 495–501
SAVERY, C. R. *see* BURY, A. F. 565–570
SCAWEN, M. & ALLEN, A. The action of proteolytic enzymes on the glycoprotein from pig gastric mucus 363–368
SCHATTENKERK, C. *see* VAN BATENBURG, O. D. 385–387
SCHMUKLER, M. *see* NEUWELT, E. A. 419–426
SELLERS, A., CARTWRIGHT, E., MURPHY, G. & REYNOLDS, J. J. Evidence that latent collagenases are enzyme–inhibitor complexes 303–307
SHIA, M. A. *see* FRIEDMAN, P. A. 39–43
SCHONFELD, S. E. *see* GUENTHER, H. L. 591–603
SIM, R. B., PORTER, R. R., REID, K. B. M. & GIGLI, I. The structure and enzymic activities of the C1r and C1s subcomponents of C1, the first component of human serum complement 219–227
SIMMONDS, R. J. & YON, R. J. Protein chromatography on adsorbents with hydrophobic and ionic groups. Purification of human erythrocyte glycophorin 397–400
SIMPSON, I. A., HOLLAWAY, M. R. & BEARD, J. The subunit structure of rabbit skeletal-muscle phosphofructokinase and the amino acid sequence of the tryptic peptide containing the highly reactive thiol group 309–316
SLAVKIN, H. C. *see* GUENTHER, H. L. 591–603
SMITH, G. D. *see* BRILEY, P. A. 325–331
SNAITH, S. M. Multiple α-mannosidase activities in mammalian tissues as shown by metal-ion activation 577–564
SOBCZAK, E. *see* STANESCU, V. 103–109
SPRAGG, S. P., ROCHE, J. J. & WILCOX, J. A comparison of the association of yeast phosphoglycerate mutase (EC 2.7.5.3) with that of haemoglobin. An ultracentrifuge study 543–555
STANESCU, V., MAROTEAUX, P. & SOBCZAK, E. Proteoglycan populations of baboon (*Papio papio*) articular cartilage. Gel-electrophoretic analysis of fractions obtained by density-gradient centrifugation and by sequential extraction 103–109
STOLL, M. S. & GRAY, C. H. The preparation and characterization of bile pigments 59–101
STRATTON, G. C. *see* GEDDES, R. 193–200
SWISHER, R. H., LANDT, M. L. & REITHEL, F. J. The molecular weight of, and evidence for two types of subunits in, the molybdenum–iron protein of *Azotobacter vinelandii* nitrogenase 427–432

TAYLAUR, C. E. *see* MILLS, G. L. 455–465
THATCHER, D. R. Enzyme instability and proteolysis during the purification of an alcohol dehydrogenase from *Drosophila melanogaster* 317–323
THOMAS, S. B. *see* GRAHAM, A. B. 117–124
TOKI, S. *see* KAGEURA, E. 401–407
TONDEUR, F. *see* BELL, M. V. 185–187
TOOGOOD, K. C. *see* GRAHAM, A. B. 117–124
TRAVIS, J. *see* JOHNSON, D. 639–641
TRENTHAM, D. R. *see* ECCLESTON, J. F. 15–29

TROWBRIDGE, I. S., NILSEN-HAMILTON, M., HAMILTON, R. T. & BEVAN, M. J. Preliminary characterization of two thymus-dependent xenoantigens from mouse lymphocytes 211–217
TROXLER, R. F. *see* BROWN, A. S. 571–581

VAN BATENBURG, O. D., VOSKUYL-HOLTKAMP, I., SCHATTENKERK, C., HOES, K., KERLING, K. E. T. & HAVINGA, E. The role of the imidazolyl nitrogen atoms of histidine-12 in ribonuclease S 385–387
VESSEY, D. A., CRISSEY, M. H. & ZAKIM, D. Kinetic studies on the enzymes conjugating bile acids with taurine and glycine in bovine liver 181–183
VESSEY, D. A. & ZAKIM, D. Characterization of microsomal choloyl-coenzyme A synthetase 357–362
VOSKUYL-HOLTKAMP, I. *see* VAN BATENBURG, O. D. 385–387

WILCOX, J. *see* SPRAGG, S. P. 543–555
WILKINSON, S. G. *see* BROWN, F. 173–175
WILLS, P. R. *see* GEDDES, R. 201–209
WINCHESTER, B. G. *see* PHILLIPS, N. C. 269–277
WOOD, G. C. *see* GRAHAM, A. B. 117–124

YON, R. J. *see* SIMMONDS, R. J. 397–400

ZAKIM, D. *see* VESSEY, D. A. 181–183, 357–362
ZALESKE, D. J. *see* DUNKER, A. K. 45–57

Index of Subjects

N-Acetylhexosaminidase, liver and colon, human, interaction of, with insolubilized concanavalin A (Brattain, M. G., Kimball, P. M., Pretlow, T. G., II & Marks, M. E.) 247–251

N-Acetylhexosaminidases, kidney, human, two, purification and properties of (Marinkovic, D. V. & Marinkovic, J. N.) 133–140

N-Acetyl-L-tryptophan, binding of bilirubin, 8-anilinonaphthalene-1-sulphonate and, to large fragments obtained by peptide digestion of human serum albumin (Geisow, M. J. & Beaven, G. H.) 477–484

Acid α-D-mannosidase, *see* α-D-Mannosidase, acid

Adenosine 3′:5′-cyclic monophosphate phosphodiesterase, baker's-yeast, binding to subcellular particles, catalytic properties and gel-filtration behaviour of (Londesborough, J.) 467–476

Adenosine triphosphatase, myosin, muscle, decapod-crustacean, regulation by calcium ions of the activity of (Lehman, W.) 291–296

Adenosine triphosphatase, myosin, skeletal-muscle, rabbit, use of chromophoric nucleotides derived from thioinosine and thioguanosine for investigation of the mechanism of the reaction catalysed by (Eccleston, J. F. & Trentham, D. R.) 15–29

Adenosine triphosphatase, sodium-plus-potassium ion-dependent, gill, eel, activation by univalent cations of (Bell, M. V., Tondeur, F. & Sargent, J. R.) 185–187

Adenosine triphosphate sulphurylase, baker's-yeast, activation of selenate by (Dilworth, G. L. & Bandurski, R. S.) 521–529

Adenylate deaminase, adenosine triphosphate-activated, body-wall-muscle, lugworm partial purification and properties of (Gibbs, K. L. & Bishop, S. H.) 511–516

Albumin, chicken-egg, acid dissociation constants of glycopeptides obtained by enzymic hydrolysis of (Austen, B. M. & Marshall, R. D.) 31–38

Albumin, serum, human, physical and binding properties of large fragments obtained by peptic digestion of (Geisow, M. J. & Beaven, G. H.) 477–484

Alcaligenes faecalis A.T.C.C. 8750, purification and properties of adenosine triphosphate-dependent deoxyribonuclease from (Rosamund, J. D. C. & Lunt, M. R.) 485–494

Alcohol dehydrogenase, fruitfly (*Drosophila melanogaster*), enzyme instability and proteolysis during the purification of (Thatcher, D. R.) 317–323

Aldehyde dehydrogenase, mitochondrial, liver, sheep, purification and properties of (Hart, G. J. & Dickinson, F. M.) 261–267

Allophycocyanin, *Cyanidium caldarium*, properties and *N*-terminal amino acid sequence of (Brown, A. S. & Troxler, R. F.) 571–581

Amelogenesis, identification of four extracellular-matrix enamel proteins during, in the rabbit embryo (Guenther, H. L., Croissant, R. D., Schonfeld, S. E. & Slavkin, H. C.) 591–603

Amino acid sequence, determination of, of locust cytochrome *c* (Lyddiatt, A. & Boulter, D.) 333–338

Amino acid sequence, determination of, of the tryptic peptide of rabbit skeletal-muscle phosphofructokinase that contains the highly active thiol group (Simpson, I. A., Hollaway, M. R. & Beard, J.) 309–316

Amino acid sequence, *N*-terminal, determination of, of *Cyanidium caldarium* allophycocyanin (Brown, A. S. & Troxler, R. F.) 571–581

Amino acid sequences, use of *NN*-dimethylaminoazobenzene 4′-isothiocyanate for high-sensitivity analysis of, of peptides and proteins (Chang, J. Y.) 517–520

Amino groups, free and bound, different reactivities of, in human deoxy- and liganded haemoglobin (Bresciani, D.) 393–395

o-Aminophenol, possible multiple binding sites for, on rat liver microsomal uridine diphosphate glucuronyltransferase (Howland, R. D. & Bohm, L. D.) 125–131

Ammonia, role of lugworm body-wall muscle adenosine triphosphate-activated adenylate deaminase in the formation of (Gibbs, K. L. & Bishop, S. H.) 511–516

Anguilla anguilla, *see* Eel

8-Anilinonaphthalene-1-sulphonate, binding of bilirubin, *N*-acetyl-L-tryptophan and, to large fragments obtained by peptic digestion of human serum albumin (Geisow, M. J. & Beaven, G. H.) 477–484

Antigens, xeno-, thymus-gland-dependent, lymphocyte, mouse, preliminary characterization of (Trowbridge, I. S., Nilsen-Hamilton, M., Hamilton, R. T. & Bevan, M. J.) 211–217

α_1-Antitrypsin, plasma, human, inactivation of, by thiol proteinases (Johnson, D. & Travis, J.) 639–641

Apomyoglobin, sperm-whale, kinetics of the recombination reaction between alkaline haematin and (Adams, P. A.) 153–158

Arenicola cristata, *see* Lugworm

Arylamidases, heart-muscle, rat, two, separation and properties of (Bury, A. F., Coolbear, T. & Savery, C. R.) 565–570

Aspartate aminotransferase, heart, pig, [^{19}F]fluorine nuclear-magnetic-resonance study of the interaction of difluoro-oxaloacetate with (Briley, P. A., Eisenthal, R., Harrison, R. & Smith, G. D.) 325–331

Aspartylglucosaminuria, activity of aspartylglucosaminase in liver, brain and kidney from normal human subjects and from patients with (Dugal, B.) 9–14

Aspartylglucosylaminase, activity of, in liver, brain and kidney from normal human subjects and from patients with aspartylglucosaminuria (Dugal, B.) 9–14

Azotobacter vinelandii A.T.C.C. 13705, molecular weight and subunit structure of the molybdenum–iron protein component of nitrogenase from (Swisher, R. H., Landt, M. L. & Reithel, F. J.) 427–432

Baboon (*Papio papio*), gel-electrophoretic analysis of proteoglycans obtained by density-gradient centrifugation and by sequential extraction from articular cartilage from (Stanescu, V., Maroteaux, P. & Sobczak, E.) 103–109

Bacteriophage λ, map of the cleavage sites for endodeoxyribonuclease *Ava*I in the chromosome of (Hughes, S. G.) 503–509

Bacteriorhodopsin, *Halobacterium halobium*, stereochemical considerations for constructing α-helical bundles with particular application to, and other membrane proteins (Dunker, A. K. & Zaleske, D. J.) 45–57

Bile acids, effects of, on the activity of guinea-pig liver microsomal choloyl-coenzyme A synthetase (Vessey, D. A. & Zakim, D.) 357–362
Bile acids, kinetic studies on the enzymes catalysing the conjugation of, with taurine and glycine in ox liver (Vessey, D. A., Crissey, M. H. & Zakim D.) 181–183
Bile pigments, preparation and characterization of (Stoll, M. S. & Gray, C. H.) 59–101
Bilirubin, binding of 8-anilinonaphthalene-1-sulphonate, *N*-acetyl-L-tryptophan and, to large fragments obtained by peptic digestion of human serum albumin (Geisow, M. J. & Beaven, G. H.) 477–484
Biliverdins, preparation and characterization of, and other bile pigments (Stoll, M. S. & Gray, C. H.) 59–101
Biliviolins, preparation and characterization of, and other bile pigments (Stoll, M. S. & Gray, C. H.) 59–101
Biotin, apparent absence of involvement of, in the vitamin K-dependent carboxylation of glutamic acid residues of proteins in rat liver microsomal fraction (Friedman, P. A. & Shia, M. A.) 39–43
Blood plasma, human, inactivation by thiol proteinases of α_1 proteinase inhibitor from (Johnson, D. & Travis, J.) 639–641
Blood plasma, human, kinetic properties of the primary plasmin inhibitor from (Christensen, U. & Clemmensen, I.) 389–391
Blood plasma, human, use of primary binding tests for the immunological characterization of ribonucleases from, and other tissues (Neuwelt, E. A., Schmukler, M., Niziak, M. S., Jewett, P. B. & Levy, C. C.) 419–426
Blood plasma, post-heparin, pig, partial purification and properties of two triacylglycerol lipases from (Ehnholm, C., Bensadoun, A. & Brown, W. V.) 347–355
Blood serum, shark, characterization of the lipoproteins of (Mills, G. L., Taylaur, C. E., Chapman, M. J. & Forster, G. R.) 455–465
Bone, rabbit, evidence that latent collagenases from, and other tissues are enzyme–inhibitor complexes (Sellers, A., Cartwright, E., Murphy, G. & Reynolds, J. J.) 303–307
Bone, rat and ox, characterization of the products of reduction of collagen after treatment of, and skin and tendon with sodium cyanoborohydride (Robins, S. P. & Bailey, A. J.) 339–346
Bradykinin, distribution and properties of peptidases acting on, in various rabbit tissues (Cicilini, M. A., Caldo, H., Berti, J. D. & Camargo, A. C. M.) 433–439
Brain, human, activity of aspartylglucosylaminase in, and other tissues from normal subjects and from patients with aspartylglucosaminuria (Dugal, B.) 9–14
Brain, rat, rapid procedure for the selective isolation of the major glycoprotein from myelin of (Quarles, R. H. & Pasnak, C. F.) 635–637
Brain cortex, ox, use of extraction with detergents and urea and of gel filtration in the preparation of fractions enriched in phosphorylated proteins from the synaptic membrane of (Dunkley, P. R., Holmes, H. & Rodnight, R.) 369–378

Calcium ions, comparison of the binding of, to the related human salivary phosphoproteins protein C and protein A (Bennick, A.) 241–245
Calcium ions, regulation by, of the adenosine triphosphatase activity of decapod-crustacean muscle myosin (Lehman, W.) 291–296
Carboxypeptidase B, pancreas, human, purification and properties of (Marinkovic, D. V., Marinkovic, J. N., Erdös, E. G. & Robinson, C. J. G.) 253–260
Cartilage, articular, baboon, gel-electrophoretic analysis of proteoglycans obtained by density-gradient centrifugation and by sequential extraction from (Stanescu, V., Maroteaux, P. & Sobczak, E.) 103–109
Catalase, liver, mouse, purified, presence of a high-molecular-weight form of the enzyme in (Baird, M. B., Massie, H. R. & Birnbaum, L. S.) 449–453
Cations, univalent, activation by, of eel gill sodium-plus-potassium ion-dependent adenosine triphosphatase (Bell, M. V., Tondeur, F. & Sargent, J. R.) 185–187
Centrophorus squamosus, *see* Shark
Cerebral cortex, ox, use of extraction with detergents and urea and of gel filtration in the preparation of fractions enriched in phosphorylated proteins from the synaptic membrane of (Dunkley, P. R., Holmes, H. & Rodnight, R.) 369–378
Chicken egg, acid dissociation constants of glycopeptides obtained by enzymic hydrolysis of albumin from (Austen, B. M. & Marshall, R. D.) 31–38
4-Chloro-7-nitrobenzofurazan, highly electrophilic character of, and possible consequences for its application as a labelling agent for proteins (Baines, B. S., Allen, G. & Brocklehurst, K.) 189–192
Chololyl-coenzyme A synthetase, microsomal, liver, guinea-pig, properties of (Vessey, D. A. & Zakim, D.) 357–362
Chromosome, bacteriophage λ, map of the cleavage sites for endodeoxyribonuclease *Ava*I in (Hughes, S. G.) 503–509
Chromous ions, reduction by, of *Pseudomonas aeruginosa* cytochrome c_{551} oxidase (Barber, D., Parr, S. R. & Greenwood, C.) 629–632
α-Chymotrypsin, use of, as a catalyst for the synthesis of peptides (Morihara, K. & Oka, T.) 531–542
Clearing-factor lipase, *see* Lipase, clearing-factor
Cobalt ions, use of activation by, and other metal ions to demonstrate multiple activities of α-mannosidase in various rat tissues (Snaith, S. M.) 557–564
Collagen, characterization of the products of reduction of, after treatment of rat and ox skin, tendon and bone with sodium cyanoborohydride (Robins, S. P. & Bailey, A. J.) 339–346
Collagen, tendon, rabbit, content of proteoglycans and the axial periodicity of (Gillard, G. C., Merrilees, M. J., Bell-Booth, P. G., Reilly, H. C. & Flint, M. H.) 145–151
Collagen fibres, dermal, human, exclusion of dextrans by meshworks of (Pearce, R. H. & Laurent, T. C.) 617–625
Collagenase inhibitor, bone, rabbit, partial characterization of (Sellers, A., Cartwright, E., Murphy, G. & Reynolds, J. J.) 303–307
Collagenases, latent, evidence that, from various rabbit tissues are enzyme–inhibitor complexes (Sellers, A., Cartwright, E., Murphy, G. & Reynolds, J. J.) 303–307

Colon, human, interaction of insolubilized concanavalin A with *N*-acetylhexosaminidase from, and other tissues (Brattain, M. G., Kimball, P. M., Pretlow, T. G., II & Marks, M. E.) 247–251
Complement, human, structure and enzymic activities of subcomponents C1r and C1s of component C1 of (Sim, R. B., Porter, R. R., Reid, K. B. M. & Gigli, I.) 219–227
Component C1, structure and enzymic activities of subcomponents C1r and C1s of, of human complement (Sim, R. B., Porter, R. R., Reid, K. B. M. & Gigli, I.) 219–227
Concanavalin A, insolubilized, interaction of, with *N*-acetylhexosaminidase from human liver and other tissues (Brattain, M. G., Kimball, P. M., Pretlow, T. G., II & Marks, M. E.) 247–251
Cyanidium caldarium, properties and *N*-terminal amino acid sequence of allophycocyanin from (Brown, A. S. & Troxler, R. F.) 571–581
Cyanoborohydride, sodium, characterization of the products of reduction of collagen after treatment of rat and ox skin, tendon and bone with (Robins, S. P. & Bailey, A. J.) 339–346
Cyclic adenosine 3′:5′-monophosphate, *see* Adenosine 3′:5′-cyclic monophosphate
Cytochrome *c*, locust, amino acid sequence of (Lyddiatt, A. & Boulter, D.) 333–338
Cytochrome c_{551} oxidase, *Pseudomonas aeruginosa*, reduction by chromous ions of (Barber, D., Parr, S. R. & Greenwood, C.) 629–632

Deoxyribonuclease, adenosine triphosphate-dependent, *Alcaligenes faecalis* A.T.C.C. 8750, purification and properties of (Rosamund, J. D. C. & Lunt, M. R.) 485–494
Deoxyribonuclease *Ava*I, endo-, map of the cleavage sites for, in the chromosome of bacteriophage λ (Hughes, S. G.) 503–509
Deoxyribonucleic acid, bacteriophage λ, map of the cleavage sites for endodeoxyribonuclease *Ava*I in (Hughes, S. G.) 503–509
Deoxyribonucleic acid, nucleosomal, rye-embryo, analysis of (Cheah, K. S. E. & Osborne, D. J.) 141–144
Dermis, human, exclusion of dextrans by meshworks of collagen fibres from (Pearce, R. H. & Laurent, T. C.) 617–625
Dextrans, exclusion of, by meshworks of human dermal collagen fibres (Pearce, R. H. & Laurent, T. C.) 617–625
Diethyl pyrocarbonate, binding of triethyltin to cat haemoglobin and modification of the binding sites by (Elliott, B. M. & Aldridge, W. N.) 583–589
Difluoro-oxaloacetate, [^{19}F]fluorine nuclear-magnetic-resonance study of the interaction of, with pig heart aspartate aminotransferase (Briley, P. A., Eisenthal, R., Harrison, R. & Smith, G. D.) 325–331
Dihydrohydroxymerodesmosine, identification of, as a product of the reduction of collagen after treatment of rat and ox skin, tendon and bone with sodium cyanoborohydride (Robins, S. P. & Bailey, A. J.) 339–346
Dihydroxyfumarate, generation of hydrogen peroxide, superoxide radical ions and hydroxy radicals during the oxidation of, by horseradish peroxidase (Halliwell, B.) 441–448
4-*NN*-Dimethylaminoazobenzene 4′-isothiocyanate, use of, for high-sensitivity analysis of amino acid sequences of peptides and proteins (Chang, J. Y.) 517–520
Diphenyleneiodonium, specific labelling by, of a constituent polypeptide of ox heart mitochondrial reduced nicotinamide–adenine dinucleotide dehydrogenase (Ragan, C. I. & Bloxham, D. P.) 605–615
Drosophila melanogaster, *see* Fruitfly

Eel (*Anguilla anguilla*), activation by univalent cations of sodium-plus-potassium ion-dependent adenosine triphosphatase from gills of (Bell, M. V., Tondeur, F. & Sargent, J. R.) 185–187
Egg, chicken, acid dissociation constants of glycopeptides obtained by enzymic hydrolysis of albumin from (Austen, B. M. & Marshall, R. D.) 31–38
Enamel proteins, extracellular-matrix, four, identification of, formed during the development of rabbit-embryo tooth organs (Guenther, H. L., Croissant, R. D., Schonfeld, S. E. & Slavkin, H. C.) 591–603
Endodeoxyribonuclease *Ava*I, map of the cleavage sites for, in the chromosome of bacteriophage λ (Hughes, S. G.) 503–509
Endopeptidases, distribution and properties of, that hydrolyse bradykinin in various rabbit tissues (Cicilini, M. A., Caldo, H., Berti, J. D. & Camargo, A. C. M.) 433–439
Enzymes, application of topological methods in kinetics of reactions catalysed by (Indge, K. J.) 633–634
Enzymes, characteristics of optimal catalysis evaluated for the simplest plausible kinetic model and their relevance to the evolution of the catalytic power of (Brocklehurst, K.) 111–116
Epoxide hydratase, liver, rat, simple method for the purification of (Knowles, R. G. & Burchell, B.) 381–383
Equilibrium data, biochemical, recommendations for the measurement and presentation of (Interunion Commission on Biothermodynamics) 1–7
Erythrocytes, human, use of chromatography on adsorbents with hydrophobic and ionic groups for the purification of glycophorin from (Simmonds, R. J. & Yon, R. J.) 397–400
Escherichia coli M.R.E. 600, dissociation of sigma factor from ribonucleic acid polymerase from (Campbell, A. M. & Lowe, P. A.) 177–179
Ethoxyformic anhydride, *see* Diethyl pyrocarbonate

Ferrous ions, use of activation by, and other metal ions to demonstrate multiple activities of α-mannosidase in various rat tissues (Snaith, S. M.) 557–564
Fibroblasts, synovial, rabbit, evidence that latent collagenases from, and other tissues are enzyme–inhibitor complexes (Sellers, A., Cartwright, E., Murphy, G. & Reynolds, J. J.) 303–307
Flavin nucleotides, use of a differential fluorimetric technique for the assay of, in mouse and rat pancreatic islets of Langerhans (Brolin, S. E. & Ågren, A.) 159–162
Fluorine isotope (^{19}F), use of, in a nuclear-magnetic-resonance study of the interaction of difluoro-oxaloacetate with pig heart aspartate aminotransferase (Briley, P. A., Eisenthal, R., Harrison, R. & Smith, G. D.) 325–331

Frog (*Rana temporaria*), partial purification and properties of liver tyrosine aminotransferase from (Ohisalo, J. J., Andersson, S. M. & Pispa, J. P.) 411–417
Fruitfly (*Drosophila melanogaster*), enzyme instability and proteolysis during the purification of alcohol dehydrogenase from (Thatcher, D. R.) 317–323

Gastric juice, human, isolation and characterization of a glycopeptide from (Goso, K. & Hotta, K.) 169–172
Gastric mucus, pig, action of proteinases on the glycoprotein from (Scawen, M. & Allen, A.) 363–368
Gills, eel, activation by univalent cations of sodium-plus-potassium ion-dependent adenosine triphosphatase from (Bell, M. V., Tondeur, F. & Sargent, J. R.) 185–187
Glutamate dehydrogenase, liver, ox, evidence from circular-dichroism studies for a conformational change in the ternary complex of, with oxidized nicotinamide–adenine dinucleotide and glutarate (Chen, S.-S., Engel, P. C. & Bayley, P. M.) 297–302
Glutamic acid residues, apparent absence of involvement of biotin in the vitamin K-dependent carboxylation of, of proteins in rat liver microsomal fraction (Friedman, P. A. & Shia, M. A.) 39–43
Glutarate, evidence from circular-dichroism studies for a conformational change in the ternary complex of ox liver glutamate dehydrogenase with oxidized nicotinamide–adenine dinucleotide and (Chen, S.-S., Engel, P. C. & Bayley, P. M.) 297–302
Glycine, kinetic studies on the enzymes catalysing the conjugation of bile acids with taurine and, in ox liver (Vessey, D. A., Crissey, M. H. & Zakim, D.) 181–183
Glycogen, influence of lysosomes on the metabolism of, in rat liver (Geddes, R. & Stratton, G. C.) 193–200
Glycogen, molecular size and shape of, from rabbit and rat liver (Geddes, R., Harvey, J. D. & Wills, P. R.) 201–209
Glycopeptide, isolation and characterization of, from human gastric juice (Goso, K. & Hotta, K.) 169–172
Glycopeptides, acid dissociation constants of, obtained by enzymic hydrolysis of chicken-egg albumin (Austen, B. M. & Marshall, R. D.) 31–38
Glycophorin, erythrocyte, human, use of chromatography on adsorbents with hydrophobic and ionic groups for the purification of (Simmonds, R. J. & Yon, R. J.) 397–400
Glycoprotein, gastric-mucus, pig, action of proteinases on (Scawen, M. & Allen, A.) 363–368
Glycoprotein, myelin, brain, rat, major, rapid procedure for the selective isolation of (Quarles, R. H. & Pasnak, C. F.) 635–637
Glycoproteins (xenoantigens), thymus-gland-dependent, lymphocyte, mouse, preliminary characterization of (Trowbridge, I. S., Nilsen-Hamilton, M., Hamilton, R. T. & Bevan, M. J.) 211–217
Glycosaminoglycans, content of, and the axial periodicity of rabbit tendon collagen (Gillard, G. C., Merrilees, M. J., Bell-Booth, P. G., Reilly, H. C. & Flint, M. H.) 145–151

Haematin, alkaline, kinetics of the recombination reaction between sperm-whale apomyoglobin and (Adams, P. A.) 153–158
Haemoglobin, cat, binding of triethyltin to, and modification of the binding sites by diethyl pyrocarbonate (Elliott, B. M. & Aldridge, W. N.) 583–589
Haemoglobin, deoxy- and liganded, human, different reactivities of free and bound amino groups in (Bresciani, D.) 393–395
Haemoglobin, human, use of ultracentrifugal analysis for comparison of the association of yeast phosphoglycerate mutase with (Spragg, S. P., Roche, J. J. & Wilcox, J.) 543–555
Halobacterium halobium, stereochemical considerations for constructing α-helical bundles with particular application to bacteriorhodopsin from, and other membrane proteins (Dunker, A. K. & Zaleske, D. J.) 45–57
Heart, ox, specific labelling by diphenyleneiodonium of a constituent polypeptide of mitochondrial reduced nicotinamide–adenine dinucleotide dehydrogenase from (Ragan, C. I. & Bloxham, D. P.) 605–615
Heart, pig, [^{19}F]fluorine nuclear-magnetic-resonance study of the interaction of difluoro-oxaloacetate with aspartate aminotransferase from (Briley, P. A., Eisenthal, R., Harrison, R. & Smith, G. D.) 325–331
Heart muscle, rat, separation and properties of two arylamidases from (Bury, A. F., Coolbear, T. & Savery, C. R.) 565–570
Hen, *see* Chicken
Hexosaminidase, see *N*-Acetylhexosaminidase
Hexosaminidases, kidney, human, two, purification and properties of (Marinkovic, D. V. & Marinkovic, J. N.) 133–140
Histidine residue 12, role of the imidazolyl nitrogen atoms of, of ribonuclease S (van Batenburg, O. D., Voskuyl-Holtkamp, I., Schattenkerk, C., Hoes, K., Kerling, K. E. T. & Havinga, E.) 385–387
Homarus americanus, *see* Lobster
Homogentisate, inhibition by, of the activity of frog liver tyrosine aminotransferase (Ohisalo, J. J., Andersson, S. M. & Pispa, J. P.) 411–417
Horseradish, generation of hydrogen peroxide, superoxide radical ions and hydroxyl radicals during the oxidation of dihydroxyfumarate by peroxidase from (Halliwell, B.) 441–448
Hydrogen peroxide, generation of superoxide radical ions, hydroxyl radicals and, during the oxidation of dihydroxyfumarate by horseradish peroxidase (Halliwell, B.) 441–448
17β-Hydroxy steroids, activity of a new guinea-pig liver testosterone 17β-dehydrogenase towards (Kageura, E. & Toki, S.) 401–407
Hydroxyl radicals, generation of hydrogen peroxide, superoxide radical ions and, during the oxidation of dihydroxyfumarate by horseradish peroxidase (Halliwell, B.) 441–448
Hydroxylases, molybdenum-containing, oxidation–reduction potentials of turkey liver xanthine dehydrogenase and the origins of oxidase and dehydrogenase behaviour in (Barber, M. J., Bray, R. C., Cammack, R. & Coughlan, M. P.) 279–289

Intestine, large, human, interaction of insolubilized concanavalin A with *N*-acetylhexosaminidase from, and other tissues (Brattain, M. G., Kimball, P. M., Pretlow, T. G., II & Marks, M. E.) 247–251

Islets of Langerhans, pancreatic, mouse and rat, use of a differential fluorimetric technique for the assay of flavin nucleotides in (Brolin, S. E. & Ågren, A.) 159–162

Kidney, human, activity of aspartylglucosylaminase in, and other tissues from normal subjects and from patients with aspartylglucosaminuria (Dugal, B.) 9–14
Kidney, human, purification and properties of two hexosaminidases from (Marinkovic, D. V. & Marinkovic, J. N.) 133–140
Kidney, ox, serological investigation of acid α-D-mannosidase from, in normal Angus cattle and in a calf with mannosidosis (Phillips, N. C., Winchester, B. G. & Jolly, R. D.) 269–277
Kidney, pig, novel system for the two-dimensional electrophoresis of membrane proteins exemplified by microvillus membrane proteins from (Booth, A. G.) 165–168
Kidney, rabbit, distribution and properties of peptidases that hydrolyse bradykinin in, and other tissues (Cicilini, M. A., Caldo, H., Berti, J. D. & Camargo, A. C. M.) 433–439
Kininase, activity of, of peptidases in various rabbit tissues (Cicilini, M. A., Caldo, H., Berti, J. D. & Camargo, A. C. M.) 433–439

Large intestine, *see* Intestine, large
Lecithin, *see* Phosphatidylcholine
Leucine 4-nitroanilide, activities of two rat heart-muscle arylamidases towards lysine 4-nitroanilide and (Bury, A. F., Coolbear, T. & Savery, C. R.) 565–570
Lipase, clearing-factor, similarity of the properties of one of two pig post-heparin plasma triacylglycerol lipases with those of (Ehnholm, C., Bensadoun, A. & Brown, W. V.) 347–355
Lipopolysaccharide, *Pseudomonas maltophilia* N.C.T.C. 10257, identification of 3-*O*-methyl-L-xylose as a component of (Brown, F., Neal, D. J. & Wilkinson, S. G.) 173–175
Lipoprotein lipase, *see* Lipase, clearing-factor
Lipoproteins, serum, shark, characterization of (Mills, G. L., Taylaur, C. E., Chapman, M. J. & Forster, G. R.) 455–465
Liver, frog, partial purification and properties of tyrosine aminotransferase from (Ohisalo, J. J., Andersson, S. M. & Pispa, J. P.) 411–417
Liver, guinea-pig, depletion of and re-activation by phospholipids of microsomal uridine diphosphate glucuronyltransferase from (Graham, A. B., Pechey, D. T., Toogood, K. C., Thomas, S. B. & Wood, G. C.) 117–124
Liver, guinea-pig, properties of microsomal chololylcoenzyme A synthetase from (Vessey, D. A. & Zakim, D.) 357–362
Liver, guinea-pig, purification and properties of a new nicotinamide–adenine dinucleotide phosphate-linked testosterone 17β-dehydrogenase from (Kageura, E. & Toki, S.) 401–407
Liver, human, activity of aspartylglucosylaminase in, and other tissues from normal subjects and from patients with aspartylglucosaminuria (Dugal, B.) 9–14
Liver, human, interaction of insolubilized concanavalin A with *N*-acetylhexosaminidase from, and other tissues (Brattain, M. G., Kimball, P. M., Pretlow, T. G., II & Marks, M. E.) 247–251
Liver, human, use of primary binding tests for the immunological characterization of ribonucleases from, and other tissues (Neuwelt, E. A., Schmukler, M., Niziak, M. S., Jewett, P. B. & Levy, C. C.) 419–426
Liver, mouse, presence of a high-molecular-weight form of the enzyme in catalase purified from (Baird, M. B., Massie, H. R. & Birnbaum, L. S.) 449–453
Liver, ox, evidence from circular-dichroism studies for a conformational change in the ternary complex of glutamate dehydrogenase from, with oxidized nicotinamide–adenine dinucleotide and glutarate (Chen, S.-S., Engel, P. C. & Bayley, P. M.) 297–302
Liver, ox, kinetic studies on the enzymes catalysing the conjugation of bile acids with taurine and glycine in (Vessey, D. A., Crissey, M. H. & Zakim, D.) 181–183
Liver, rabbit, and rat, molecular size and shape of glycogen from (Geddes, R., Harvey, J. D. & Wills, P. R.) 201–209
Liver, rat, apparent absence of involvement of biotin in the vitamin K-dependent carboxylation of glutamic residues of proteins in microsomal fraction from (Friedman, P. A. & Shia, M. A.) 39–43
Liver, rat, influence of lysosomes on the metabolism of glycogen in (Geddes, R. & Stratton, G. C.) 193–200
Liver, rat, possible multiple binding sites for *o*-aminophenol on microsomal uridine diphosphate glucuronyltransferase from (Howland, R. D. & Bohm, L. D.) 125–131
Liver, rat, simple method for the purification of epoxide hydratase from (Knowles, R. G. & Burchell, B.) 381–383
Liver, rat, use of activation by metal ions to demonstrate multiple activities of α-mannosidase in, and other tissues (Snaith, S. M.) 557–564
Liver, sheep, purification and properties of mitochondrial aldehyde dehydrogenase from (Hart, G. J. & Dickinson, F. M.) 261–267
Liver, turkey, oxidation–reduction potentials of xanthine dehydrogenase from, and the origins of oxidase and dehydrogenase behaviour in molybdenum-containing hydroxylases (Barber, M. J., Bray, R. C., Cammack, R. & Coughlan, M. P.) 279–289
Lobster (*Homarus americanus*), regulation by calcium ions of the adenosine triphosphatase activity of myosin from muscle from, and other decapod crustaceans (Lehman, W.) 291–296
Locust (*Schistocerca gregaria*), amino acid sequence of cytochrome *c* from (Lyddiatt, A. & Boulter, D.) 333–338
Lugworm (*Arenicola cristata*), partial purification and properties of adenosine triphosphate-activated adenylate deaminase from the body-wall muscle of (Gibbs, K. L. & Bishop, S. H.) 511–516
Lung, rabbit, distribution and properties of peptidases that hydrolyse bradykinin in, and other tissues (Cicilini, M. A., Caldo, H., Berti, J. D. & Camargo, A. C. M.) 433–439

Lymphocytes, mouse, preliminary characterization of two thymus-gland-dependent xenoantigens from (Trowbridge, I. S., Nilsen-Hamilton, M., Hamilton, R. T. & Bevan, M. J.) 211–217
Lysine 4-nitroanilide, activities of two rat heart-muscle arylamidases towards leucine 4-nitroanilide and (Bury, A. F., Coolbear, T. & Savery, C. R.) 565–570
Lysosomes, liver, rat, influence of, on the metabolism of glycogen (Geddes, R. & Stratton, G. C.) 193–200

Manganese ions, use of activation by, and other metal ions to demonstrate multiple activities of α-mannosidase in various rat tissues (Snaith, S. M.) 557–564
α-D-Mannosidase, acid, kidney, ox, serological investigation of, in normal Angus cattle and in a calf with mannosidosis (Phillips, N. C., Winchester, B. G. & Jolly, R. D.) 269–277
α-Mannosidase, liver and testis, rat, use of activation by metal ions to demonstrate multiple activities of (Snaith, S. M.) 557–564
Mannosidosis, serological investigation of kidney acid α-D-mannosidase in normal Angus cattle and in a calf with (Phillips, N. C., Winchester, B. G. & Jolly, R. D.) 269–277
Membrane, erythrocyte, human, use of chromatography on adsorbents with hydrophobic and ionic groups for the purification of glycophorin from (Simmonds, R. J. & Yon, R. J.) 397–400
Membrane, synaptic, cerebral-cortex, ox, use of extraction with detergents and urea and of gel filtration in the preparation of fractions enriched in phosphorylated proteins from (Dunkley, P. R., Holmes, H. & Rodnight, R.) 369–378
Membrane proteins, novel system for the two-dimensional electrophoresis of (Booth, A. G.) 165–168
Membranes, baker's-yeast, reversible binding of adenosine 3′:5′-cyclic monophosphate phosphodiesterase to (Londesborough, J.) 467–476
Membranes, microsomal, liver, guinea-pig, depletion of and re-activation by phospholipids of uridine diphosphate glucuronyltransferase of (Graham, A. B., Pechey, D. T., Toogood, K. C., Thomas, S. B. & Wood, G. C.) 117–124
Membranes, stereochemical considerations for constructing α-helical protein bundles with particular application to proteins of (Dunker, A. K. & Zaleske, D. J.) 45–57
3-*O*-Methyl-L-xylose, identification of, as a component of *Pseudomonas maltophilia* N.C.T.C. 10257 lipopolysaccharide (Brown, F., Neal, D. J. & Wilkinson, S. G.) 173–175
Microsomal fraction, liver, guinea-pig, depletion of and re-activation by phospholipids of uridine diphosphate glucuronyltransferase of (Graham, A. B., Pechey, D. T., Toogood, K. C., Thomas, S. B. & Wood, G. C.) 117–124
Microsomal fraction, liver, guinea-pig, properties of chololyl-coenzyme A synthetase from (Vessey, D. A. & Zakim, D.) 357–362
Microsomal fraction, liver, rat, apparent absence of involvement of biotin in the vitamin K-dependent carboxylation of glutamic acid residues of proteins in (Friedman, P. A. & Shia, M. A.) 39–43
Microsomal fraction, liver, rat, possible multiple binding sites for *o*-aminophenol on, uridine diphosphate glucuronyltransferase from (Howland, R. D. & Bohm, L. D.) 125–131
Mitochondria, heart, ox, specific labelling by diphenyleneiodonium of a constituent polypeptide of reduced nicotinamide–adenine dinucleotide dehydrogenase from (Ragan, C. I. & Bloxham, D. P.) 605–615
Mitochondria, liver, sheep, purification and properties of aldehyde dehydrogenase from (Hart, G. J. & Dickinson, F. M.) 261–267
Molybdenum, oxidation–reduction potentials of turkey liver xanthine dehydrogenase and the origins of oxidase and dehydrogenase behaviour in hydroxylases containing (Barber, M. J., Bray, R. C., Cammack, R. & Coughlan, M. P.) 279–289
Molybdenum–iron protein, nitrogenase, *Azotobacter vinelandii* A.T.C.C. 13705, molecular weight and subunit structure of (Swisher, R. H., Landt, M. L. & Reithel, F. J.) 427–432
Mucus, gastric, pig, action of proteinases on the glycoprotein from (Scawen, M. & Allen, A.) 363–368
Muscle, body-wall, lugworm, partial purification and properties of adenosine triphosphate-activated adenylate deaminase from (Gibbs, K. L. & Bishop, S. H.) 511–516
Muscle, decapod-crustacean, regulation by calcium ions of the adenosine triphosphatase activity of myosin from (Lehman, W.) 291–296
Muscle, heart, rat, separation and properties of two arylamidases from (Bury, A. F., Coolbear, T. & Savery, C. R.) 565–570
Muscle, skeletal, rabbit, interaction of chromophoric nucleotides with subfragment 1 of myosin from (Eccleston, J. F. & Trentham, D. R.) 15–29
Muscle, skeletal, rabbit, subunit structure of phosphofructokinase from, and the amino acid sequence of the tryptic peptide containing the highly active thiol group (Simpson, I. A., Hollaway, M. R., & Beard, J.) 309–316
Myelin, brain, rat, rapid procedure for the selective isolation of the major glycoprotein from (Quarles, R. H. & Pasnak, C. F.) 635–637
Myoglobin, apo-, sperm-whale, kinetics of the recombination reaction between alkaline haematin and (Adams, P. A.) 153–158
Myosin, muscle, decapod-crustacean, regulation by calcium ions of the adenosine triphosphatase activity of (Lehman, W.) 291–296
Myosin, skeletal-muscle, rabbit, interaction of chromophoric nucleotides with subfragment 1 of (Eccleston, J. F. & Trentham, D. R.) 15–29

Nicotinamide–adenine dinucleotide, oxidized, evidence from circular-dichroism studies for a conformational change in the ternary complex of ox liver glutamate dehydrogenase with glutarate and (Chen, S.-S., Engel, P. C. & Bayley, P. M.) 297–302
Nicotinamide–adenine dinucleotide (reduced) dehydrogenase, mitochondrial, heart, ox, specific labelling by diphenyleneiodonium of a constituent polypeptide of (Ragan, C. I. & Bloxham, D. P.) 605–615

Nitrogenase, *Azotobacter vinelandii* A.T.C.C. 13705, molecular weight and subunit structure of the molybdenum–iron protein component of (Swisher, R. H., Landt, M. L. & Reithel, F. J.) 427–432
Nucleosomal fraction, rye-embryo, analysis of deoxyribonucleic acid from (Cheah, K. S. E. & Osborne, D. J.) 141–144
Nucleotides, chromophoric, interaction of, derived from thioinosine and thioguanosine with subfragment 1 of rabbit skeletal-muscle myosin (Eccleston, J. F. & Trentham, D. R.) 15–29

Oligonucleotides, separation by high-voltage electrophoresis on diethylaminoethyl-paper and by thin-layer chromatography of, obtained by ribonuclease digestion of baker's-yeast valine transfer ribonucleic acid 2b (Gorbulev, V. G., Kutateladze, T. V., Barciszewski, J. & Axel'rod, V. D.) 409–410
Ovalbumin, chicken-egg, acid dissociation constants of glycopeptides obtained by enzymic hydrolysis of (Austen, B. M. & Marshall, R. D.) 31–38

Pancreas, human, purification and properties of carboxypeptidase B from (Marinkovic, D. V., Marinkovic, J. N., Erdös, E. G. & Robinson, C. J. G.) 253–260
Pancreas, human, use of primary binding tests for the immunological characterization of ribonucleases from, and other tissues (Neuwelt, E. A., Schmukler, M., Niziak, M. S., Jewett, P. B. & Levy, C. C.) 419–426
Pancreas, mouse and rat, use of a differential fluorimetric technique for the assay of flavin nucleotides in islets of Langerhans of (Brolin, S. E. & Ågren, A.) 159–162
Papio papio, *see* Baboon
Parotid gland, human, binding of calcium ions to the phosphoprotein protein C of saliva secreted by, in comparison with the binding of calcium ions to the related phosphoprotein protein A (Bennick, A.) 241–245
Parotid gland, human, chemical and physical characterization of the phosphoprotein protein C of saliva secreted by, and comparison with the related phosphoprotein protein A (Bennick, A.) 229–239
Pepsin, action of, and other proteinases on the glycoprotein from pig gastric mucus (Scawen, M. & Allen, A.) 363–368
Peptidase (carboxypeptidase B), pancreas, human, purification and properties of (Marinkovic, D. V., Marinkovic, J. N., Erdös, E. G. & Robinson, C. J. G.) 253–260
Peptidases, distribution and properties of, that hydrolyse bradykinin in various rabbit tissues (Cicilini, M. A., Caldo, H., Berti, J. D. & Camargo, A. C. M.) 433–439
Peptides, use of α-chymotrypsin as a catalyst for the synthesis of (Morihara, K. & Oka, T.) 531–542
Peptides, use of 4-*NN*-dimethylaminoazobenzene 4′-isothiocyanate for high-sensitivity analysis of amino acid sequences of proteins and (Chang, J. Y.) 517–520
Peroxidase, horseradish, generation of hydrogen peroxide, superoxide radical ions and hydroxyl radicals during the oxidation of dihydroxyfumarate by (Halliwell, B.) 441–448
Phage, *see* Bacteriophage
Phosphatidylcholine, depletion of and re-activation by, and other phospholipids of guinea-pig liver microsomal uridine diphosphate glucuronyltransferase (Graham, A. B., Pechey, D. T., Toogood, K. C., Thomas, S. B. & Wood, G. C.) 117–124
Phosphofructokinase, skeletal-muscle, rabbit, subunit structure of, and the amino acid sequence of the tryptic peptide containing the highly active thiol group (Simpson, I. A., Hollaway, M. R. & Beard, J.) 309–316
Phosphoglycerate mutase, yeast, use of ultracentrifugal analysis for comparison of the association of, with that of human haemoglobin (Spragg, S. P., Roche, J. J. & Wilcox, J.) 543–555
Phospholipids, depletion of and re-activation by, of guinea-pig liver microsomal uridine diphosphate glucuronyltransferase (Graham, A. B., Pechey, D. T., Toogood, K. C., Thomas, S. B. & Wood, G. C.) 117–124
Phosphoprotein (protein C), salivary, human, binding of calcium ions to, in comparison with the binding of calcium ions to the related phosphoprotein protein A (Bennick, A.) 241–245
Phosphoprotein (protein C), salivary, human, chemical and physical characterization of, and comparison with the related phosphoprotein protein A (Bennick, A.) 229–239
Phycocyanin, allo-, *Cyanidium caldarium*, properties and *N*-terminal amino acid sequence of (Brown, A. S. & Troxler, R. F.) 571–581
Plasma, human, inactivation by thiol proteinases of α_1 proteinase inhibitor from (Johnson, D. & Travis, J.) 639–641
Plasma, human, kinetic properties of the primary plasmin inhibitor from (Christensen, U. & Clemmensen, I.) 389–391
Plasma, human, use of primary binding tests for the immunological characterization of ribonucleases from, and other tissues (Neuwelt, E. A., Schmukler, M., Niziak, M. S., Jewett, P. B. & Levy, C. C.) 419–426
Plasma, post-heparin, pig, partial purification and properties of two triacylglycerol lipases from (Ehnholm, C., Bensadoun, A. & Brown, W. V.) 347–355
Plasmin inhibitor, plasma, human, primary, kinetic properties of (Christensen, U. & Clemmensen, I.) 389–391
Polypeptides, use of α-chymotrypsin as a catalyst for the synthesis of (Morihara, K. & Oka, T.) 531–542
Pronase, action of, and other proteinases on the glycoprotein from pig gastric mucus (Scawen, M. & Allen, A.) 363–368
Protease, *see* Proteinase
Protein, molybdenum–iron, nitrogenase, *Azotobacter vinelandii* A.T.C.C. 13705, molecular weight and subunit structure of (Swisher, R. H., Landt, M. L. & Reithel, F. J.) 427–432
Protein A, salivary, human, comparison of the binding of calcium ions to, and protein C (Bennick, A.) 241–245
Protein A, salivary, human, comparison of the chemical and physical properties of, and protein C (Bennick, A.) 229–239
Protein C, salivary, human, binding of calcium ions to, in comparison with the binding of calcium ions to the related protein A (Bennick, A.) 241–245

Protein C, salivary, human, chemical and physical characterization of, and comparison with the related protein A (Bennick, A) 229–239
Protein–polysaccharides, *see* Proteoglycans
α_1 Proteinase inhibitor, plasma, human, inactivation of, by thiol proteinases (Johnson, D. & Travis, J.) 639–641
Proteinases, action of, on the glycoprotein from pig gastric mucus (Scawen, M. & Allen, A.) 363–368
Proteinases, thiol, inactivation by, of human plasma α_1 proteinase inhibitor (Johnson, D. & Travis, J.) 639–641
Proteins, apparent absence of involvement of biotin in the vitamin K-dependent carboxylation of glutamic acid residues of, in rat liver microsomal fraction (Friedman, P. A. & Shia, M. A.) 39–43
Proteins, enamel, extracellular-matrix, four, identification of, formed during the development of rabbit-embryo tooth organs (Guenther, H. L., Croissant, R. D., Schonfeld, S. E. & Slavkin, H. C.) 591–603
Proteins, highly electrophilic character of 4-chloro-7-nitrobenzofurazan and possible consequences for its application as a labelling agent for (Baines, B. S., Allen, G. & Brocklehurst, K.) 189–192
Proteins, membrane, novel system for the two-dimensional electrophoresis of (Booth, A. G.) 165–168
Proteins, membrane, stereochemical considerations for constructing α-helical protein bundles with particular application to (Dunker, A. K. & Zaleske, D. J.) 45–57
Proteins, phosphorylated, use of extraction with detergents and urea and of gel filtration in the preparation of fractions enriched in, from ox cerebral-cortex synaptic membrane (Dunkley, P. R., Holmes, H. & Rodnight, R.) 369–378
Proteins, use of *NN*-dimenthylaminoazobenzene 4′-isothiocyanate from high-sensitivity analysis of amino acid sequences of peptides and (Chang, J. Y.) 517–520
Proteoglycans, articular-cartilage, baboon, gel-electrophoretic analysis of, obtained by density-gradient centrifugation and by sequential extraction (Stanescu, V., Maroteaux, P. & Sobczak, E.) 103–109
Proteoglycans, content of, and the axial periodicity of rabbit tendon collagen (Gillard, G. C., Merrilees, M. J., Bell-Booth, P. G., Reilly, H. C. & Flint, M. H.) 145–151
Pseudomonas aeruginosa, reduction by chromous ions of cytochrome c_{551} oxidase from (Barber, D., Parr, S. R. & Greenwood, C.) 629–632
Pseudomonas maltophilia N.C.T.C. 10257, identification of 3-*O*-methyl-L-xylose as a component of the lipopolysaccharide of (Brown, F., Neal, D. J. & Wilkinson, S. G.) 173–175

Rana temporaria, *see* Frog
Red blood cells, *see* Erythrocytes
Ribonuclease S, role of the imidazolyl nitrogen atoms of histidine residue 12 of (van Batenburg, O. D., Voskuyl-Holtcamp, I., Schattenkerk, C., Hoes, K., Kerling, K. E. T. & Havinga, E.) 385–387
Ribonucleases, plasma, pancreas, spleen and liver, human, use of primary binding tests for the immunological characterization of (Neuwelt, E. A., Schmukler, M., Niziak, M. S., Jewett, P. B. & Levy, C. C.) 419–426
Ribonucleic acid polymerase, *Escherichia coli* M.R.E. 600, dissociation of sigma factor from (Campbell, A. M. & Lowe, P. A.) 177–179
Ribonucleic acid, transfer, valine, 2b, baker's-yeast, separation by high-voltage electrophoresis on diethylaminoethyl-paper and by thin-layer chromatography of oligonucleotides obtained by ribonuclease digestion of (Gorbulev, V. G., Kutateladze, T. V., Barciszewski, J. & Axel'rod, V. D.) 409–410
Rumen, ox, purification and properties of urease from (Mahadevan, S., Sauer, F. D. & Erfle, J. D.) 495–501
Rye (*Secale cereale*) embryos, analysis of nucleosomal deoxyribonucleic acid from (Cheah, K. S. E. & Osborne, D. J.) 141–144

Saccharomyces cerevisiae, activation of selenate by adenosine triphosphate sulphurylase from (Dilworth, G. L. & Bandurski, R. S.) 521–529
Saccharomyces cerevisiae, binding to subcellular particles, catalytic properties and gel-filtration behaviour of adenosine 3′:5′-cyclic monophosphate phosphodiesterase from (Londesborough, J.) 467–476
Saccharomyces cerevisiae, separation by high-voltage electrophoresis on diethylaminoethyl-paper and by thin-layer chromatography of oligonucleotides obtained by ribonuclease digestion of valine transfer ribonucleic acid 2b from (Gorbulev, V. G., Kutateladze, T. V., Barciszewski, J. & Axel'rod, V. D.) 409–410
Saliva, human, binding of calcium ions to the phosphoprotein protein C of, in comparison with the binding of calcium ions to the related phosphoprotein protein A (Bennick, A.) 241–245
Saliva, human, chemical and physical characterization of the phosphoprotein protein C of, and comparison with the related phosphoprotein protein A (Bennick, A) 229–239
Salivary gland, parotid, human, binding of calcium ions to the phosphoprotein protein C of saliva secreted by, in comparison with the binding of calcium ions to the related phosphoprotein protein A (Bennick, A.) 241–245
Salivary gland, parotid, human, chemical and physical characterization of the phosphoprotein protein C of saliva secreted by, and comparison with the related phosphoprotein protein A (Bennick, A.) 229–239
Schistocerca gregaria, *see* Locust
Secale cereale, *see* Rye
Selenate, activation of, by baker's-yeast adenosine triphosphate sulphurylase (Dilworth, G. L. & Bandurski, R. S.) 521–529
Serum, human, structure and enzymic activities of subcomponents C1r and C1s of component C1 of complement from (Sim, R. B., Porter, R. R., Reid, K. B. M. & Gigli, I.) 219–227
Serum albumin, *see* Albumin, serum
Serum lipoproteins, *see* Lipoproteins, serum
Shark (*Centrophorus squamosus*), characterization of the serum lipoproteins of (Mills, G. L., Taylaur, C. E., Chapman, M. J. & Forster, G. R.) 455–465
Sigma factor, dissociation of, from *Escherichia coli* M.R.E. 600 ribonucleic acid polymerase (Campbell, A. M. & Lowe, P. A.) 177–179
Skeletal muscle, *see* Muscle, skeletal

Skin, human, exclusion of dextrans by meshworks of collagen fibres from (Pearce, R. H. & Laurent, T. C.) 617–625
Skin, rabbit, evidence that latent collagenases from, and other tissues are enzyme–inhibitor complexes (Sellers, A., Cartwright, E., Murphy, G. & Reynolds, J. J.) 303–307
Skin, rat and ox, characterization of the products of reduction of collagen after treatment of, and bone and tendon with sodium cyanoborohydride (Robins, S. P. & Bailey, A. J.) 339–346
Sodium cyanoborohydride, characterization of the products of reduction of collagen after treatment of rat and ox skin, tendon and bone with (Robins, S. P. & Bailey, A. J.) 339–346
Sperm whale, kinetics of the recombination reaction between alkaline haematin and apomyoglobin from (Adams, P. A.) 153–158
Spleen, human, use of primary binding tests for the immunological characterization of ribonucleases from, and other tissues (Neuwelt, E. A., Schmukler, M., Niziak, M. S., Jewett, P. B. & Levy, C. C.) 419–426
Steroids, 17β-hydroxy, activity of a new guinea-pig liver testosterone 17β-dehydrogenase towards (Kageura, E. & Toki, S.) 401–407
Stomach, human, isolation and characterization of a glycopeptide from gastric juice from (Goso, K. & Hotta, K.) 169–172
Stomach, pig, action of proteinases on the glycoprotein from gastric mucus from (Scawen, M. & Allen, A.) 363–368
Subcomponents C1r and C1s, structure and enzymic activities of, of component C1 of human complement (Sim, R. B., Porter, R. R., Reid, K. B. M. & Gigli, I.) 219–227
Sulphydryl, *see* Thiol
Superoxide radical ions, generation of hydrogen peroxide, hydroxyl radicals and, during the oxidation of dihydroxyfumarate by horseradish peroxidase (Halliwell, B.) 441–448
Synaptic membrane, *see* Membrane, synaptic

Taurine, kinetic studies on the enzymes catalysing the conjugation of bile acids with glycine and, in ox liver (Vessey, D. A., Crissey, M. H. & Zakim, D.) 181–183
Tendon, rabbit, content of proteoglycans and the axial periodicity of collagen from (Gillard, G. C., Merrilees, M. J., Bell-Booth, P. G., Reilly, H. C. & Flint, M. H.) 145–151
Tendon, rat and ox, characterization of the products of reduction of collagen after treatment of, and bone and skin with sodium cyanoborohydride (Robins, S. P. & Bailey, A. J.) 339–346
Testis, rat, use of activation by metal ions to demonstrate multiple activities of α-mannosidase in, and other tissues (Snaith, S. M.) 557–564
Testosterone 17β-dehydrogenase, nicotinamide–adenine dinucleotide phosphate-linked, liver, guinea-pig, new, purification and properties of (Kageura, E. & Toki, S.) 401–407
Thioguanosine, interaction of chromophoric nucleotides derived from thioinosine and, with subfragment 1 of rabbit skeletal-muscle myosin (Eccleston, J. F. & Trentham, D. R.) 15–29
Thioinosine, interaction of chromophoric nucleotides derived from thioguanosine and, with subfragment 1 of rabbit skeletal-muscle myosin (Eccleston, J. F. & Trentham, D. R.) 15–29
Thiol group, highly active, amino acid sequence of the tryptic peptide of rabbit skeletal-muscle phosphofructokinase that contains (Simpson, I. A., Hollaway, M. R. & Beard, J.) 309–316
Tooth organs, rabbit-embryo, identification of four extracellular-matrix enamel proteins formed during the development of (Guenther, H. L., Croissant, R. D., Schonfeld, S. E. & Slavkin, H. C.) 591–603
Transfer ribonucleic acid, *see* Ribonucleic acid, transfer
Triacylglycerol lipases, plasma, post-heparin, pig, two, partial purification and properties of (Ehnholm, C., Bensadoun, A. & Brown, W. V.) 347–355
Triethyltin, binding of, to cat haemoglobin and modification of the binding sites by diethyl pyrocarbonate (Elliott, B. M. & Aldridge, W. N.) 583–589
Triglyceride, *see* Triacylglycerol
Trypsin, action of, and other proteinases on the glycoprotein from pig gastric mucosa (Scawen, M. & Allen, A.) 363–368
Turkey, oxidation–reduction potentials of liver xanthine dehydrogenase from and the origins of oxidase and dehydrogenase behaviour in molybdenum-containing hydroxylases (Barber, M. J., Bray, R. C., Cammack, R. & Coughlan, M. P.) 279–289
Tyrosine aminotransferase, liver, frog, partial purification and properties of (Ohisalo, J. J., Andersson, S. M. & Pispa, J. P.) 411–417

Urease, rumen, ox, purification and properties of (Mahadevan, S., Sauer, F. D. & Erfle, J. D.) 495–501
Uridine diphosphate glucuronyltransferase, microsomal, liver, guinea-pig, depletion of and re-activation by phospholipids of (Graham, A. B., Pechey, D. T., Toogood, K. C., Thomas, S. B. & Wood, G. C.) 117–124
Uridine diphosphate glucuronyltransferase, microsomal, liver, rat, possible multiple binding sites for *o*-aminophenol on (Howland, R. D. & Bohm, L. D.) 125–131
Uterus, rabbit, evidence that latent collagenases from, and other tissues are enzyme–inhibitor complexes (Sellers, A., Cartwright, E., Murphy, G. & Reynolds, J. J.) 303–307

Valine transfer ribonucleic acid, baker's-yeast, separation by high-voltage electrophoresis on diethylaminoethyl-paper and by thin-layer chromatography of oligonucleotides obtained by ribonuclease digestion of (Gorbulev, V. G., Kutateladze, T. V., Barciszewski, J. & Axel'rod, V. D.) 409–410
Vitamin K, apparent absence of involvement of biotin in the carboxylation of glutamic acid residues of rat liver microsomal proteins that is dependent on (Friedman, P. A. & Shia, M. A.) 39–43

Xanthine dehydrogenase, liver, turkey, oxidation–reduction potentials of, and the origins of oxidase and dehydrogenase behaviour in molybdenum-containing hydroxylases (Barber, M. J., Bray, R. C., Cammack, R. & Coughlan, M. P.) 279–289

Xenoantigens, thymus-gland-dependent, lymphocyte, mouse, preliminary characterization of (Trowbridge, I. S., Nilsen-Hamilton, M., Hamilton, R. T. & Bevan, M. J.) 211–217

Yeast, baker's (*Saccharomyces cerevisiae*), activation of selenate by adenosine triphosphate sulphurylase from (Dilworth, G. L. & Bandurski, R. S.) 521–529

Yeast, baker's (*Saccharomyces cerevisiae*), binding to subcellular particles, catalytic properties and gel-filtration behaviour of adenosine 3′:5′-cyclic monophosphate phosphodiesterase from (Londesborough, J.) 467–476

Yeast, baker's (*Saccharomyces cerevisiae*), separation by high-voltage electrophoresis on diethylaminoethyl-paper and by thin-layer chromatography of oligonucleotides obtained by ribonuclease digestion of valine transfer ribonucleic acid 2b from (Gorbulev, V. G., Kutateladze, T. V., Barciszewski, J. & Axel'rod, V. D.) 409–410

Yeast, use of ultracentrifugal analysis for comparison of the association of phosphoglycerate mutase from, with that of human haemoglobin (Spragg, S. P., Roche, J. J. & Wilcox, J.) 543–555

ECTS

tion by
oethyl-
oligo-
ion of
rbulev,
xel'rod,

rison of
from,
Roche,

1977